

Last Time

Greedy Flip Algorithm takes $O(n^2)$ time.

Any Δ^n can be flipped to Delaunay

Today Randomized Incremental Construction


... but first, yet another definition of
(characterization)
the Delaunay Δ^n .

Claim: Among all Δ^n s of a point set P , the Delaunay Δ^n maximizes the minimum angle.

pf idea: Show that flipping an edge that is not LD can only increase the smallest angle.

So, our algorithmic result, gives a way to prove facts about Delp.


First, a refresher from high school/ancient Greece.



Thales Thm If \overline{ab} is the diameter of circle (a, b, c) , then $\angle acb$ is 90° .


pf $2\alpha + 2\beta = 180^\circ$
 $\Rightarrow \angle acb = \alpha + \beta = 90^\circ$.

This is true more generally.



If $\triangle abc$ and $\triangle abc'$ have the same circumcircle, then $\angle acb = \angle ac'b$.


pf




$2\alpha + 2\beta + 2\gamma = 180^\circ$
 $\Rightarrow \angle acb = \alpha + \beta = 90 - \gamma$

For c' , α and β change but γ is the same so $\alpha + \beta$ does not change. **Be careful: Add signed angles.**

So,





$$\theta_1 > \theta_4$$

$$\theta_2 > \theta_7$$

$$\theta_6 > \theta_3$$

$$\theta_5 > \theta_8$$

So, every angle after the flip is bigger than some angle that appeared before the flip.

Incremental Algorithms


A staple of computational geometry

Idea: Add points one at a time.

Invariant: After i points added, we have computed the Del. Δ^n of those i pts.

1st Step Start with convex hull.

But for now assume its a triangle.



2nd Step Add one new point.

- Find Δ containing p_{i+1}
- Split it into 3
- Run the Greedy Flip Algorithm.



3rd Step Repeat.

Question: How many flips?


Some observations

- 1) All new triangles ^{in step i} have a vertex at p_i .
- 2) Every flip adds one edge incident to p_i .
- 3) No flip removes an edge incident to p_i .

\Rightarrow degree(p_i) - 3 flips in step i .

\uparrow This is the degree
in $\text{Del}\{p_1, \dots, p_i\}$.

This could still be quadratic.



First $n/2$ points "near" a circle.
Next $n/2$ points all require
 $n/2 - 2$ flips.

\Rightarrow $\Theta(n^2)$ total flips.

Idea: Add the points in a random order

all permutations have equal prob.

Let $\langle p_1, \dots, p_n \rangle$ be the random order.

Let $Q_i = \{p_1, \dots, p_i\}$ (Q_i is the first i pts in the order)

We want to know the degree of p_i in $\text{Del}(Q_i)$,
~~be~~ call it δ_i .

~~we~~ We know $\# \text{flips} < \sum_{i=1}^n \delta_i$

What about $E[\# \text{flips}]$?

Expectation over all random orderings.

Use linearity of expectations:

$$E[\# \text{flips}] < E\left[\sum_{i=1}^n \delta_i\right] = \sum_{i=1}^n E[\delta_i]$$